

The Future of Sample Preparation:
Scaling Medical Device, Biopharma, and Reference Lab Workflows

FerroTec marcus evans

TUES, 29TH SEPT, 2026 LIVE WEBINAR

The views expressed during this webinar belong solely to the participants based on their experience as a business professional. These views do not represent those of the participants' employers or other group(s) they may be affiliated with and they are not an endorsement of any solutions or services associated with a webinar including those of the Marcus Evans Group.

The Future of Sample Preparation: Scaling Medical Device, Biopharma, and Reference Lab Workflows



DR. TAKKI Q. MOMIN
President & Medical Director
Clinical Strategy & Diagnostics
Piedmont Labs



DR. JEANINE VAN NOSTRAND
Assistant Professor
Department of Molecular
and Cellular Biology
Baylor College of Medicine



DR. SADAF KHAN
General Manager
Life Sciences Division
Ferrotec

High-throughput molecular diagnostics, next-generation sequencing, and translational research place heavy demands on clinical laboratories. Healthcare systems and diagnostic manufacturers invest substantial capital into advanced detection instruments. These platforms process large batches of samples quickly. However, laboratory throughput depends on a stage that receives far less attention: pre-analytical sample preparation.

Sample preparation remains the primary source of manual labor, technical error, and protocol variance. Whether extracting nucleic acids for clinical monitoring or isolating proteins for oncology research, analytical reliability depends on upstream specimen handling. Laboratory executives, clinical directors, and research heads are recognizing that downstream detection platforms cannot compensate for poor initial sample quality.

Achieving scalable growth requires modernizing sample preparation workflows through automated platforms and integrated digital systems.

The True Cost of Pre-Analytical Inefficiencies

Workflow friction in both clinical laboratories and translational research centers originates primarily in upstream processing. When initial handling fails, the resulting errors create compounding operational costs.


In translational research, these inefficiencies often result in inconclusive experiments. Minor pipetting variations, technical mistakes during cell lysis, and uneven protein loading during Western blot analysis generate experimental noise. When data is inconclusive, research teams must discard results and rerun the protocol. Repeating these multi-step experiments consumes expensive reagents, wastes laboratory hours, and stalls project milestones.

The consequences are even more severe in high-complexity clinical diagnostics. Clinical workflows operate under strict turnaround time windows. Patient care pathways depend on prompt, accurate test reports. A typical clinical specimen moves through multiple distinct operational stages:

- **Processing:** Technicians perform accessioning, sample lysis, and target isolation.
- **Interpretation:** Analysts review amplification curves, sequencing reads, and quality metrics.
- **Validation:** Laboratory directors verify internal controls and release finalized clinical reports.



Because a specimen passes through several hands, operational variance accumulates at each stage. Biological specimens such as whole blood, tissue biopsies, and viral swabs are susceptible to degradation. Variable shipping conditions and repeated freeze-thaw cycles compromise nucleic acid stability. When a clinical sample degrades, or an extraction step fails quality control, the laboratory faces an unwanted choice. It must either restart a 24- to 48-hour testing cycle or request a recollection from the patient. In acute care, oncology testing, and infectious disease screening, these delays disrupt patient management and place heavy operational strain on referring clinics.



Eliminating Bottlenecks: Upstream Quality in Automated Systems

A disciplined operational principle in clinical laboratory management is to trace every workflow bottleneck to its root cause. When an assay fails, managers should avoid making superficial adjustments. Instead, they need to map the entire process from collection to reporting. This systematic evaluation determines whether an error stems from mechanical equipment failure, facility environmental variations, or protocol design flaws.

System mapping highlights a major risk in automated testing environments: high-speed automation amplifies the impact of poor input quality. Modern analyzers process thousands of reactions rapidly. However, feeding compromised, degraded, or contaminated samples into automated systems simply generates invalid data faster. Producing unreliable results at high volume creates significant financial waste and clinical risk.

To address this challenge, automated sample preparation systems focus on condensing complex, multi-step chemistries into fast, reliable isolation protocols. Traditional extraction methods relied on lengthy manual protocols, including repeated centrifugation, toxic organic solvents, and delicate phase separations. Newer extraction platforms utilize specialized magnetic and ferrofluid materials, a technology originally developed for aerospace engineering. These systems capture total nucleic acids through simplified, single-tube or automated-cartridge workflows.

By completing isolation protocols in 15 to 25 minutes, modern platforms minimize manual touchpoints and reduce liquid transfer errors. Operational simplicity is just as important as processing speed:

- Effective automation platforms avoid complex hardware interfaces that require weeks of vendor training.
- Standardized, self-contained instruments allow technicians to run protocols with minimal onboarding.
- Straightforward interfaces minimize user error and protect operations against high personnel turnover.

Maximizing Human Capital: The Real Return on Automation

Budget proposals for automated laboratory instrumentation often focus on capital purchase costs and consumable pricing. Experienced laboratory directors look at a more valuable metric: the strategic deployment of human capital.

Molecular biologists, certified clinical laboratory scientists, and doctoral researchers possess specialized training in data analysis, experimental design, and clinical correlation. Yet traditional laboratory workflows force these skilled professionals to spend hours on repetitive, manual benchwork. Technicians spend significant portions of their shifts pipetting 96-well plates, running routine verification gels, and monitoring extraction columns.

Automating sample extraction returns two to three hours of productive time to each technician every day. Reallocating this labor provides immediate organizational benefits:

- **Accelerated Diagnostics:** Clinical personnel direct their time toward assay development, internal quality audits, and resolving data discrepancies.
- **Expanded Research Capacity:** Translational scientists spend less time on routine bench tasks and more time designing experiments, analyzing complex datasets, and testing new research avenues.
- **Interdisciplinary Collaboration:** Reducing manual bench burdens allows wet-lab processors to coordinate directly with bioinformaticians, pathologists, and clinical teams.




The return on automation is not measured merely in labor hours saved. It is measured in the increased scientific output and diagnostic reliability of the existing workforce.

Operational Reproducibility: Unifying Standardization and Flexibility

Laboratory leaders frequently face a difficult operational balance: enforcing strict protocol standardization while preserving scientific flexibility. Clinical diagnostics and translational discovery address this challenge from different perspectives, yet both require reliable sample preparation.

In clinical diagnostics, reproducibility requires strict adherence to Standard Operating Procedures and comprehensive digital tracking. Quality control systems must record every process variable, including sample transport temperatures, reagent lot numbers, and instrument runtime logs.

Far from restricting innovation, rigorous standardization actually protects it. When baseline processing remains uniform, unexpected assay variations stand out immediately as true biological signals rather than manual pipetting errors.



For instance, laboratories operating in regions with severe seasonal heat and humidity often experience analyte degradation during sample transit. Because baseline protocols are standardized, laboratory teams can review thousands of data points to pinpoint stability loss caused by thermal shifts. This insight allows scientists to make targeted chemical adjustments, such as adding stabilizing phosphate buffers to preserve analytes during summer freeze-thaw cycles. Without standardized baselines, identifying such specific environmental variables amid technical noise is virtually impossible.

In translational research, scientists work with unknown biological systems. In this setting, reproducible experimental controls are vital. If control samples fluctuate due to manual handling differences, researchers can spend months investigating artificial phenotypes. Scientific flexibility belongs in the research questions being asked, such as evaluating novel biomarkers or adjusting treatment timepoints. The underlying sample preparation protocol must remain rigid and automated.

Maintaining Regulatory Compliance Across Growing Operations

As diagnostic laboratories expand their testing menus, regulatory compliance grows more demanding. Operations must satisfy overlapping standards established by federal agencies, state departments of health, and independent accreditation organizations such as CLIA, CAP, and COLA.

These regulatory frameworks safeguard clinical testing quality. However, compiling compliance documentation across high testing volumes creates heavy administrative burdens. Manual recordkeeping and subjective sample assessments leave laboratories vulnerable during inspections.

Automated sample preparation and digital tracking systems serve as an essential compliance framework:

- **Digital Audit Trails:** Automated platforms log cycle durations, reagent volumes, and processing parameters, creating complete records for every processed sample.
- **Variance Reduction:** Eliminating manual steps prevents technicians from taking unauthorized protocol shortcuts during high-volume shifts.
- **Defensible Inspection Data:** Integrated systems produce objective, reproducible run records that simplify quarterly reporting and satisfy multi-agency audits.

The Implementation Framework: People, Process, and Technology

Integrating automated sample preparation into an established laboratory requires a balanced operational strategy. To prevent costly equipment purchases from sitting idle, leadership teams should align implementation around three key areas:



- **People:** Build cultural trust across the technical staff before installation. Select user-friendly systems that require minimal training so technicians embrace the technology quickly. Shift the organizational mindset from manual sample manipulation toward process oversight.

- **Process:** Map the specimen journey from accessioning to downstream analysis to eliminate hidden handoff delays. Establish universal extraction protocols that support multiple diagnostic tests across shared laboratory equipment.

- **Technology:** Deploy bi-directional data connections between automation platforms, Laboratory Information Management Systems, and Electronic Medical Records. Invest in flexible platforms that yield high recovery across varied sample matrices.

Laboratories should avoid purchasing single-purpose instruments that support only one project. True operational scalability comes from selecting versatile platforms that handle diverse assays over multi-year lifecycles. Sharing automated platforms across departments breaks down institutional silos and maximizes capital utilization.



Key Takeaways

- **Pre-Analytical Steps Determine Overall Velocity:** Upstream sample extraction remains the primary operational bottleneck in molecular testing. Advanced downstream sequencers and amplification platforms cannot correct for degraded, variable starting material.
- **Automation Protects Downstream Data Quality:** Modern analyzers process specimens at high speeds, which means poor input produces faulty data faster. Automating extraction eliminates pre-analytical variability and protects downstream investments.
- **Standardization Unlocks Scientific Discovery:** Enforcing rigid sample preparation protocols provides the baseline stability needed to isolate true biological discoveries and identify environmental anomalies.
- **The Core Value Is Human Capital Optimization:** The primary return on investment from automation comes from shifting skilled personnel away from repetitive bench tasks and into assay development, data interpretation, and clinical validation.
- **Sustainable Growth Requires Strategic Alignment:** Lasting operational efficiency demands the coordinated integration of People, Process, and Technology, supported by automated data exchange across clinical and research teams.